

Epidemiologic Risk Classification for Asymptomatic Healthcare Personnel Following Exposure to Patients with COVID-19 or their Secretions/Excretions in a Healthcare Setting, and their Associated Monitoring and Work Restriction Recommendations

CDC Guidance: <https://www.cdc.gov/coronavirus/2019-ncov/hcp/guidance-risk-assesment-hcp.html#table1>

HCP=healthcare personnel; PPE=personal protective equipment

Epidemiologic risk factors	Exposure category	Recommended Monitoring for COVID-19 (until 14 days after last potential exposure)	Work Restrictions for Asymptomatic HCP
Prolonged close contact with a COVID-19 patient who was wearing a facemask (i.e., source control)			
HCP PPE: None	Medium	Active	Exclude from work for 14 days after last exposure
HCP PPE: Not wearing a facemask or respirator	Medium	Active	Exclude from work for 14 days after last exposure
HCP PPE: Not wearing eye protection	Low	Self with delegated supervision	None
HCP PPE: Not wearing gown or gloves ^a	Low	Self with delegated supervision	None
HCP PPE: Wearing all recommended PPE (except wearing a facemask instead of a respirator)	Low	Self with delegated supervision	None
Prolonged close contact with a COVID-19 patient who was not wearing a facemask (i.e., no source control)			
HCP PPE: None	High	Active	Exclude from work for 14 days after last exposure
HCP PPE: Not wearing a facemask or respirator	High	Active	Exclude from work for 14 days after last exposure
HCP PPE: Not wearing eye protection ^b	Medium	Active	Exclude from work for 14 days after last exposure
HCP PPE: Not wearing gown or gloves ^{a,b}	Low	Self with delegated supervision	None
HCP PPE: Wearing all recommended PPE (except wearing a facemask instead of a respirator) ^b	Low	Self with delegated supervision	None

^aThe risk category for these rows would be elevated by one level if HCP had extensive body contact with the patients (e.g., rolling the patient).

^bThe risk category for these rows would be elevated by one level if HCP performed or were present for a procedure likely to generate higher concentrations of respiratory secretions or aerosols (e.g., cardiopulmonary resuscitation, intubation, extubation, bronchoscopy, nebulizer therapy, sputum induction). For example, HCP who were wearing a gown, gloves, eye protection and a facemask (instead of a respirator) during an aerosol-generating procedure would be considered to have a medium-risk exposure.